

NEW SHARP JORDAN TYPE INEQUALITIES AND THEIR APPLICATIONS

ZHEN-HANG YANG

ABSTRACT. In this paper, we prove that for $x \in (0, \pi/2)$

$$(\cos p_0 x)^{1/p_0} < \frac{\sin x}{x} < \left(\cos \frac{x}{3}\right)^3$$

with the best constants $p_0 = 0.347307245464\dots$ and $1/3$. Moreover, if $p \in (0, 1/3]$ then the double inequality

$$\beta_p (\cos px)^{1/p} < \frac{\sin x}{x} < (\cos px)^{1/p}$$

holds for $x \in (0, \pi/2)$, where $\beta_p = 2\pi^{-1} (\cos \frac{p\pi}{2})^{-1/p}$ and 1 are the best possible. Its reverse one holds if $p \in [1/2, 1]$. As applications, some new inequalities are established.

1. INTRODUCTION

The classical Jordan's inequality [10] states that for $x \in (0, \pi/2]$

$$(1.1) \quad \frac{2}{\pi} \leq \frac{\sin x}{x} \leq 1,$$

with equality holds if and only if $x = \pi/2$.

Some new developments on refinements, generalizations and applications of Jordan's inequality can be found in [14] and related references therein.

In [15], Qi et al. proved that

$$(1.2) \quad \cos^2 \frac{x}{2} < \frac{\sin x}{x}$$

holds for $x \in (0, \pi/2)$. Recently, Klén et al. [8, Theorem 2.4] showed that the function $p \mapsto (\cos px)^{1/p}$ is decreasing on $(0, 1)$ and proved that for $x \in (-\sqrt{27/5}, \sqrt{27/5})$

$$(1.3) \quad \cos^2 \frac{x}{2} \leq \frac{\sin x}{x} \leq \cos^3 \frac{x}{3} \leq \frac{2 + \cos x}{3}$$

hold.

This paper is motivated by these studies and it is aimed at giving the sharp bounds $(\cos px)^{1/p}$ for $(\sin x)/x$, that is, for $x \in (0, \pi/2)$, determine the best $p, q \in$

Date: April 10, 2012.

2010 Mathematics Subject Classification. 26D05, 26D15, 33F05, 26E60.

Key words and phrases. Jordan type inequalities, trigonometric functions, sharp bound, relative error.

This paper is in final form and no version of it will be submitted for publication elsewhere.

$(0, 1]$ such that

$$(1.4) \quad (\cos px)^{1/p} < \frac{\sin x}{x} < (\cos qx)^{1/q}$$

hold, respectively.

Our main results are contained the following two theorems.

Theorem 1. *Let $p, q \in (0, 1]$. Then the inequalities*

$$(1.5) \quad (\cos px)^{1/p} < \frac{\sin x}{x} < (\cos qx)^{1/q}$$

hold for $x \in (0, \pi/2)$ if and only if $p \in [p_0, 1]$ and $q \in (0, 1/3]$, where $p_0 = 0.347307245464\dots$

Moreover, we have

$$(1.6) \quad \left(\cos \frac{x}{3}\right)^\alpha < \frac{\sin x}{x} < \left(\cos \frac{x}{3}\right)^3,$$

$$(1.7) \quad (\cos p_0 x)^{1/p_0} < \frac{\sin x}{x} < (\cos p_0 x)^{1/(3p_0^2)},$$

where the exponents $\alpha = \frac{2(\ln \pi - \ln 2)}{\ln 4 - \ln 3}$, 3 , $1/p_0$, $1/(3p_0^2)$ are the best constants.

Theorem 2. *For $p \in (0, 1]$, let f_p be the function defined on $(0, \pi/2)$ by*

$$(1.8) \quad f_p(x) = \ln \frac{\sin x}{x} - \frac{1}{p} \ln \cos px.$$

Then f_p is decreasing if $p \in (0, 1/3]$ and increasing if $p \in [1/2, 1]$.

Moreover, if $p \in (0, 1/3]$ then

$$(1.9) \quad \beta_p\left(\frac{\pi}{4}\right) (\cos px)^{1/p} < \frac{\sin x}{x} < (\cos px)^{1/p}$$

with the best possible constants $\beta_p(\pi/2) = 2\pi^{-1} (\cos \frac{p\pi}{2})^{-1/p}$ and 1. (1.9) is reversed if $p \in [1/2, 1]$.

2. LEMMAS

Lemma 1 ([17], [1]). *Let $f, g : [a, b] \mapsto \mathbb{R}$ be two continuous functions which are differentiable on (a, b) . Further, let $g' \neq 0$ on (a, b) . If f'/g' is increasing (or decreasing) on (a, b) , then so are the functions*

$$x \mapsto \frac{f(x) - f(b)}{g(x) - g(b)} \quad \text{and} \quad x \mapsto \frac{f(x) - f(a)}{g(x) - g(a)}.$$

Lemma 2 ([2]). *Let a_n and b_n ($n = 0, 1, 2, \dots$) be real numbers and let the power series $A(t) = \sum_{n=1}^{\infty} a_n t^n$ and $B(t) = \sum_{n=1}^{\infty} b_n t^n$ be convergent for $|t| < R$. If $b_n > 0$ for $n = 0, 1, 2, \dots$, and a_n/b_n is strictly increasing (or decreasing) for $n = 0, 1, 2, \dots$, then the function $A(t)/B(t)$ is strictly increasing (or decreasing) on $(0, R)$.*

Lemma 3 ([7, pp.227-229]). *We have*

$$(2.1) \quad \cot x = \frac{1}{x} - \sum_{n=1}^{\infty} \frac{2^{2n}}{(2n)!} |B_{2n}| x^{2n-1}, \quad |x| < \pi,$$

$$(2.2) \quad \tan x = \sum_{n=1}^{\infty} \frac{2^{2n}-1}{(2n)!} 2^{2n} |B_{2n}| x^{2n-1}, \quad |x| < \pi/2,$$

where B_n is the Bernoulli numbers.

Lemma 4. Let F_p be the function defined $(0, \pi/2)$ by

$$(2.3) \quad F_p(x) = \frac{\ln \frac{\sin x}{x}}{\ln \cos px}.$$

Then F_p is strictly increasing on $(0, \pi/2)$ if $p \in (0, \sqrt{5}/5]$ and decreasing on $(0, \pi/2)$ if $p \in [1/2, 1]$. Moreover, we have

$$(2.4) \quad \frac{\ln 2 - \ln \pi}{\ln(\cos \frac{1}{2}\pi p)} \ln \cos px < \ln \frac{\sin x}{x} < \frac{1}{3p^2} \ln \cos px$$

if $p \in (0, \sqrt{5}/5]$. The inequalities (2.4) are reversed if $p \in [1/2, 1]$.

Proof. For $x \in (0, \pi/2)$, we define $f(x) = \ln \frac{\sin x}{x}$ and $g(x) = \ln \cos px$, where $p \in (0, 1]$. Note that $f(0^+) = g(0^+) = 0$, then $F_p(x)$ can be written as

$$F_p(x) = \frac{f(x) - f(0^+)}{g(x) - g(0^+)}.$$

Differentiation and using (2.1) and (2.2) yield

$$\frac{f'(x)}{g'(x)} = \frac{p \left(\frac{1}{x} - \cot x \right)}{\tan px} = \frac{\sum_{n=1}^{\infty} \frac{2^{2n}}{(2n)!} |B_{2n}| x^{2n-1}}{\sum_{n=1}^{\infty} \frac{2^{2n}-1}{(2n)!} p^{2n-2} 2^{2n} |B_{2n}| x^{2n-1}} := \frac{\sum_{n=1}^{\infty} a_n x^{2n-1}}{\sum_{n=1}^{\infty} b_n x^{2n-1}},$$

where

$$a_n = \frac{2^{2n}}{(2n)!} |B_{2n}|, \quad b_n = \frac{2^{2n}-1}{(2n)!} p^{2n-2} 2^{2n} |B_{2n}|.$$

Clearly, if the monotonicity of a_n/b_n is proved, then by Lemma 2 it is deduced the monotonicity of f'/g' , and then the monotonicity of the function F_p easily follows from Lemma 1. Now we prove the monotonicity of a_n/b_n . Indeed, elementary computation yields

$$\begin{aligned} \frac{b_{n+1}}{a_{n+1}} - \frac{b_n}{a_n} &= (2^{2n+2} - 1) p^{2n} - (2^{2n} - 1) p^{2n-2} \\ &= (4^{n+1} - 1) p^{2n-2} \left(p^2 - \frac{1}{4} + \frac{3}{4(4^{n+1} - 1)} \right), \end{aligned}$$

from which it is easy to obtain that for $n \in \mathbb{N}$

$$\frac{b_{n+1}}{a_{n+1}} - \frac{b_n}{a_n} \begin{cases} \leq 0 & \text{if } p^2 < \frac{1}{5}, \\ > & \text{if } p^2 \geq \frac{1}{4}. \end{cases}$$

It is seen that b_n/a_n is decreasing if $0 < p \leq \sqrt{5}/5$ and increasing if $1/2 \leq p \leq 1$, which together with $a_n, b_n > 0$ for $n \in \mathbb{N}$ leads to a_n/b_n is strictly increasing if $0 < p \leq \sqrt{5}/5$ and decreasing if $1/2 \leq p \leq 1$.

By the monotonicity of the function F_p and notice that

$$F_p(0^+) = \frac{1}{3p^2} \quad \text{and} \quad F_p\left(\frac{\pi}{2}\right) = \frac{\ln 2 - \ln \pi}{\ln(\cos \frac{1}{2}\pi p)},$$

the inequalities (2.4) follow immediately. $\square$

Lemma 5. For $p \in (0, 1]$, let f_p be the function defined on $(0, \pi/2)$ by 1.8).

(i) If $f_p(x) < 0$ holds for all $x \in (0, \pi/2)$ then $p \in (0, 1/3]$.

(ii) If $f_p(x) > 0$ for all $x \in (0, \pi/2)$, then $p \in [p_0, 1]$, where $p_0 = 0.347307245464\dots$ is the unique root of equation

$$(2.5) \quad f_p\left(\frac{\pi}{2}\right) = \ln \frac{2}{\pi} - \frac{1}{p} \ln \left(\cos \frac{1}{2}\pi p \right) = 0$$

on $(0, 1]$.

Proof. Firstly, we show that there is a unique $p_0 \in (0, 1)$ to satisfy (2.5) such that $f_p(\frac{\pi}{2}) < 0$ for $p \in (0, p_0)$ and $f_p(\frac{\pi}{2}) > 0$ for $p \in (p_0, 1]$. Indeed, as mentioned previous (see [8, Theorem 2.3]), the function $p \mapsto p^{-1} \ln(\cos \frac{1}{2}\pi p)$ is decreasing on $(0, 1)$, and therefore $p \mapsto f_p(\frac{\pi}{2})$ is increasing on $(0, 1)$. Since

$$\begin{aligned} f_{1/3}(\frac{\pi}{2}) &= \ln \frac{2}{\pi} - 3 \ln \frac{\sqrt{3}}{2} < 0, \\ f_{1/2}(\frac{\pi}{2}) &= \ln \frac{2}{\pi} - 2 \ln \frac{\sqrt{2}}{2} > 0, \end{aligned}$$

so the equation ((2.5) has a unique root p_0 on $(0, 1)$ and $p_0 \in (1/3, 1/2)$ such that $f_p(\frac{\pi}{2}) < 0$ for $p \in (0, p_0)$ and $f_p(\frac{\pi}{2}) > 0$ for $p \in (p_0, 1]$. Numerical calculation reveals that $p_0 = 0.347307245464\dots$

Secondly, if inequality $f_p(x) < 0$ holds for $x \in (0, \pi/2)$, then we have

$$\begin{cases} \lim_{x \rightarrow 0^+} \frac{f_p(x)}{x^2} = \lim_{x \rightarrow 0^+} \frac{\ln \frac{\sin x}{x} - \frac{1}{p} \ln \cos px}{x^2} = \frac{1}{2}p - \frac{1}{6} \leq 0, \\ f_p(\frac{\pi}{2}) = \ln \frac{2}{\pi} - \frac{1}{p} \ln(\cos \frac{1}{2}\pi p) \leq 0. \end{cases}$$

Solving the inequalities for p yields

$$p \in (0, 1/3] \cap (0, p_0] = (0, 1/3].$$

In the same way, if inequality $f_p(x) > 0$ for all $x \in (0, \pi/2)$, then

$$p \in [1/3, 1] \cap [p_0, 1] = [p_0, 1],$$

which completes the proof. $\square$

3. PROOFS OF MAIN RESULTS

Proof of Theorem 1. (i) We first prove the second inequality of (1.5) holds if and only if $q \in (0, 1/3]$.

The necessity follows from Lemma 5. It remains to show that the condition $q \in (0, 1/3]$ is sufficient. Since $q \in (0, 1/3] \subset (0, \sqrt{5}/5]$, by the second inequality of (2.4) it is obtained that

$$\ln \frac{\sin x}{x} < \frac{1}{3q^2} \ln \cos qx = \frac{1}{3q} \ln(\cos qx)^{1/q} \leq \ln(\cos qx)^{1/q},$$

which proves the sufficiency.

(ii) Now we show that the first inequality of (1.5) holds if and only if $p \in [p_0, 1]$.

The necessity is due to Lemma 5. We prove the first inequality of (1.5) holds if $p \in [p_0, 1]$. In fact, in view of $p_0 \in (0, \sqrt{5}/5]$ it follows from the first inequality of (2.4) that

$$\ln \frac{\sin x}{x} > \frac{\ln 2 - \ln \pi}{\ln(\cos \frac{1}{2}\pi p_0)} \ln \cos p_0 x = \frac{p_0(\ln 2 - \ln \pi)}{\ln(\cos \frac{1}{2}\pi p_0)} \ln(\cos p_0 x)^{1/p_0} = \ln(\cos p_0 x)^{1/p_0},$$

where the last equality holds is due to p_0 is the unique root of equation (2.5). And, by the monotonicity of the function $p \rightarrow (\cos px)^{1/p}$, we conclude that for $p \in [p_0, 1]$

$$\frac{\sin x}{x} > (\cos p_0 x)^{1/p_0} \geq (\cos px)^{1/p},$$

which implies the sufficiency.

(iii) Lastly, put $p = 1/3$ and p_0 in (2.4) lead to (1.6) and (1.7), respectively. $\square$

Proof of Theorem 2. Differentiation and using (2.1) and (2.2) yield

$$\begin{aligned}
f'_p(x) &= \left(\cot x - \frac{1}{x}\right) + \tan px \\
&= -\sum_{n=1}^{\infty} \frac{2^{2n}}{(2n)!} |B_{2n}| x^{2n-1} + \sum_{n=1}^{\infty} \frac{2^{2n}-1}{(2n)!} p^{2n-1} 2^{2n} |B_{2n}| x^{2n-1} \\
&= \sum_{n=1}^{\infty} \frac{(2^{2n}-1) 2^{2n}}{(2n)!} |B_{2n}| \left(p^{2n-1} - \frac{1}{2^{2n}-1}\right) x^{2n-1} \\
&: = \sum_{n=2}^{\infty} c_n d_n x^{2n-1},
\end{aligned}$$

where

$$\begin{aligned}
c_n &= \frac{(2^{2n}-1) 2^{2n}}{(2n)!} |B_{2n}| \frac{p^{2n-1} - \frac{1}{2^{2n}-1}}{p - \left(\frac{1}{2^{2n}-1}\right)^{1/(2n-1)}} > 0, \\
d_n &= p - h(n), \quad h(n) = \left(\frac{1}{2^{2n}-1}\right)^{1/(2n-1)}
\end{aligned}$$

for $n \geq 1$ and $p \in (0, 1]$.

Considering the function $g : (1/2, \infty) \mapsto (0, \infty)$ defined by

$$(3.1) \quad g(x) = \left(\frac{1}{2^{2x}-1}\right)^{1/(2x-1)},$$

and differentiation leads to

$$\begin{aligned}
\frac{2(2x-1)^2}{g(x)} g'(x) &= \ln(2^{2x}-1) - \frac{(2x-1) 2^{2x} \ln 2}{(2^{2x}-1)} := g_1(x), \\
g'_1(x) &= \frac{2^{2x+1} \ln^2 2}{(2^{2x}-1)^2} (2x-1) > 0,
\end{aligned}$$

which reveals that g_1 is increasing on $(1/2, \infty)$, and therefore $g_1(x) > g_1(1/2^+) = 0$. It follows that $g'(x) > 0$, then, g is increasing on $(1/2, \infty)$, hence for $n \geq 1$

$$\frac{1}{3} = g(1) \leq g(n) \leq g(\infty) = \lim_{n \rightarrow \infty} \left(\frac{1}{2^{2n}-1}\right)^{1/(2n-1)} = \frac{1}{2},$$

and then, $d_n = p - g(n) \leq 0$ if $p \in (0, 1/3]$ and $d_n = p - g(n) \geq 0$ if $p \in [1/2, 1]$. Thus, if $p \in (0, 1/3]$ then $f'_p(x) < 0$, that is, f_p is decreasing, and it is derived that

$$\ln \beta_p\left(\frac{\pi}{2}\right) = f_p\left(\frac{\pi}{2}^-\right) < f_p(x) < \lim_{x \rightarrow 0^+} f_p(x) = 0,$$

which yields (1.9).

Likewise, if $p \in [1/2, 1]$ then $f'_p(x) > 0$, then, f_p is increasing, and (1.9) is reversed, which completes the proof. $\square$

4. COROLLARIES

Putting $p = 1/3$ in Theorem 2, we have

Corollary 1. (i) For $x \in (0, \pi/2)$

$$(4.1) \quad \beta_{1/3}\left(\frac{\pi}{2}\right) \cos^3 \frac{x}{3} < \frac{\sin x}{x} < \cos^3 \frac{x}{3}$$

with the best constants $\beta_{1/3}(\pi/2) = 16\sqrt{3}/(9\pi) = 0.98014\dots$ and 1.

(ii) For $x \in (0, \pi/4)$ the inequalities

$$(4.2) \quad \beta_{1/3}^{(1)}\left(\frac{\pi}{4}\right) \cos^3 \frac{x}{3} < \frac{\sin x}{x} < \cos^3 \frac{x}{3}$$

hold, where $\beta_{1/3}(\pi/4) = 16(3\sqrt{3} - 5)/\pi = 0.99900\dots$ and 1 are the best possible.

Remark 1. The second inequality of (4.1) holds for $x \in (0, 3\pi/2)$. In fact, we define

$$h(x) := x - \left(\cos \frac{x}{3}\right)^{-3} \sin x$$

and differentiation yields

$$h'(x) = 1 - \frac{\sin x}{\cos^4 \frac{x}{3}} \sin \frac{x}{3} - \frac{\cos x}{\cos^3 \frac{x}{3}} = \tan^4 \frac{x}{3} > 0.$$

Hence for $x \in (0, 3\pi/2)$ then $h(x) > h(0) = 0$, which implies that the second inequality of (4.1) holds for $x \in (0, 3\pi/2)$.

Thus, by replacing x for $3x$ the second inequality of (4.1) and next using duplication formula for sine function we have

Corollary 2. For $x \in (0, \pi/2)$

$$(4.3) \quad \frac{\sin x}{x} (4 \cos^2 x - 1) = \frac{\sin 3x}{x} < 3 \cos^3 x$$

holds.

Utilizing the second inequality of (4.1) holds for $x \in (0, 3\pi/2)$ we also obtain

Corollary 3. For $x \in (0, \pi/2)$ we have

$$(4.4) \quad \frac{\sin x}{x} < \frac{3}{4} \frac{(\cos x + 1)^2}{2 \cos x + 1}.$$

Proof. Since the second inequality of (4.1) holds for $x \in (0, 3\pi/2)$, we have

$$\frac{\sin(3x/2)}{3x/2} < \cos^3 \frac{x}{2}.$$

Multiplying both sides by $\cos(x/2)$ and next using "product into sum" and half-angle formulas give us

$$(4.5) \quad \frac{\sin 2x + \sin x}{3x} < \left(\frac{\cos x + 1}{2}\right)^2,$$

which, after applying double-angle formula and next dividing both sides by $(2 \cos x + 1)/3$, is the desired inequality. $\square$

Putting $p = 1/2$ in Theorem 2, we get

Corollary 4. For $x \in (0, \pi/2)$, the inequalities

$$(4.6) \quad \left(\cos \frac{x}{2}\right)^2 < \frac{\sin x}{x} < \beta_{1/2}\left(\frac{\pi}{2}\right) \left(\cos \frac{x}{2}\right)^2$$

hold, where $\beta_{1/2}(\pi/2) = 4/\pi$ and 1 are the best possible constants.

Remark 2. The first inequality of (4.6) or (1.2) is equivalent to $\tan(x/2) > x/2$, which holds for $x/2 \in (0, \pi/2)$, that is, $x \in (0, \pi)$. The second one of (4.1) or (1.3) holds for $x \in (0, 3\pi/2)$ due to Remark previous. While the third one of (1.3) holds for $x \in (-\infty, \infty)$, since

$$\cos^3 \frac{x}{3} - \frac{2 + \cos x}{3} = -\frac{1}{3} \left(\cos \frac{x}{3} + 2 \right) \left(\cos \frac{x}{3} - 1 \right)^2 < 0.$$

Consequently, the value range of variable x such that (1.3) holds can be extended to $(0, \pi)$, which slightly improves (1.3).

5. APPLICATIONS

As consequences of main results, we will establish some new inequalities in this section. The following is a direct corollary of Theorem 2.

Corollary 5. We have

$$(5.1) \quad \left(\frac{2}{\pi} \right)^p \frac{1}{p} \tan \frac{p\pi}{2} < \int_0^{\pi/2} \left(\frac{\sin x}{x} \right)^p dx < \int_0^{\pi/2} (\cos px) = \frac{1}{p} \sin \frac{p\pi}{2}$$

if $p \in (0, 1/3]$. Inequalities (5.1) is reversed if $p \in [1/2, 1]$.

For the estimate for the sine integral defined by

$$\text{Si}(x) = \int_0^x \frac{\sin t}{t} dt,$$

there has some results, for example, Qi [13] showed that

$$1.3333\dots = \frac{4}{3} < \text{Si}\left(\frac{\pi}{2}\right) < \frac{\pi+1}{3} = 1.3805\dots;$$

the following two estimations are due to Wu [18], [19]:

$$\begin{aligned} 1.3569\dots &= \frac{\pi+5}{6} < \text{Si}\left(\frac{\pi}{2}\right) < \frac{\pi+1}{3} = 1.3805\dots, \\ 1.3688\dots &= \frac{92-\pi^2}{60} < \text{Si}\left(\frac{\pi}{2}\right) < \frac{8+4\pi}{15} = 1.3711\dots \end{aligned}$$

Now we give a more better one.

Corollary 6. We have

$$(5.2) \quad 1.3696\dots = \frac{(3\sqrt{3}-5)(2\pi+9\sqrt{3}+22)}{2\pi} < \text{Si}\left(\frac{\pi}{2}\right) < \frac{2\pi+9\sqrt{3}+22}{32} = 1.3710\dots$$

Proof. By 2, for $x \in (0, \pi/2)$ we have

$$\beta_{1/3}\left(\frac{\pi}{4}\right) \cos^3 \frac{x}{6} < \frac{\sin \frac{x}{2}}{\frac{x}{2}} < \cos^3 \frac{x}{6},$$

where $\beta_{1/3}(\pi/4) = 16(3\sqrt{3}-5)/\pi$, and multiplying both sides by $\cos(x/2)$ leads to

$$(5.3) \quad \beta_{1/3}\left(\frac{\pi}{4}\right) \cos^3 \frac{x}{6} \cos \frac{x}{2} < \frac{\sin x}{x} = \frac{\sin \frac{x}{2}}{\frac{x}{2}} \cos \frac{x}{2} < \cos^3 \frac{x}{6} \cos \frac{x}{2}.$$

Integrating both sides over $[0, \pi/2]$ yields

$$\beta_{1/3}\left(\frac{\pi}{4}\right) \int_0^{\pi/2} \cos^3 \frac{x}{6} \cos \frac{x}{2} dx < \int_0^{\pi/2} \frac{\sin x}{x} dx < \int_0^{\pi/2} \cos^3 \frac{x}{6} \cos \frac{x}{2} dx.$$

From the following

$$\int_0^{\pi/2} \cos \frac{x}{2} \cos^3 \frac{x}{6} dx = \frac{2\pi + 9\sqrt{3} + 22}{32}$$

(5.1) follows. □

The Catalan constant [5]

$$K = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^2} = 0.9159655941772190\dots$$

is a famous mysterious constant appearing in many places in mathematics and physics. Its integral representations contain the following [3]

$$K = \int_0^1 \frac{\arctan x}{x} dx = \frac{1}{2} \int_0^{\pi/2} \frac{x}{\sin x} dx.$$

We present two estimations for K below.

Corollary 7. *We have*

$$(5.4) \quad \frac{1}{8} (4 + 3 \ln 3) < K < \frac{3\sqrt{3}\pi}{128} (4 + 3 \ln 3),$$

$$(5.5) \quad \frac{8\sqrt{3}}{9} \ln(1 + \sqrt{3}) - \frac{40 - 22\sqrt{3}}{3} < K < \pi \left(\frac{9 + 5\sqrt{3}}{36} \ln(\sqrt{3} + 1) - \frac{1 + 5\sqrt{3}}{48} \right).$$

Proof. By inequalities (4.1) we get for $(0, \pi/2)$

$$\frac{1}{\cos^3 \frac{x}{3}} < \frac{x}{\sin x} < \frac{9\pi}{16\sqrt{3}} \frac{1}{\cos^3 \frac{x}{3}},$$

and integrating both sides over $[0, \pi/2]$ yields

$$\int_0^{\pi/2} \frac{dx}{\cos^3 \frac{x}{3}} < \int_0^{\pi/2} \frac{x}{\sin x} dx < \int_0^{\pi/2} \frac{dx}{\cos^3 \frac{x}{3}}.$$

From the

$$\int_0^{\pi/2} \frac{dx}{\cos^3 \frac{x}{3}} = 1 + \frac{3 \ln 3}{4}$$

(5.4) follows.

Now we prove (5.5). From (5.3) it is deduced that

$$\frac{1}{\cos \frac{x}{2} \cos^3 \frac{x}{6}} < \frac{x}{\sin x} < \frac{1}{\beta_{1/3}(\pi/4)} \frac{1}{\cos \frac{x}{2} \cos^3 \frac{x}{6}},$$

and integrating over $[0, \pi/2]$ yields

$$\int_0^{\pi/2} \frac{dx}{\cos \frac{x}{2} \cos^3 \frac{x}{6}} < \int_0^{\pi/2} \frac{x}{\sin x} dx < \frac{1}{\beta_{1/3}(\pi/4)} \int_0^{\pi/2} \frac{dx}{\cos \frac{x}{2} \cos^3 \frac{x}{6}}.$$

Direct computation gives

$$\begin{aligned} \int \frac{dt}{\cos 3t \cos^3 t} &= \frac{\sin t}{576 \cos^5 t} - \frac{559 \sin t}{1152 \cos^3 t} - \frac{113 \sin t}{384 \cos t} + \frac{65 \sin^3 t}{192 \cos^3 t} + \frac{37 \sin^5 t}{1152 \cos^5 t} \\ &\quad - \frac{13 \sin^5 t}{384 \cos^5 t} + \frac{4\sqrt{3}}{27} \ln \frac{1 + \sin(2t - \frac{\pi}{6})}{1 - \sin(2t + \frac{\pi}{6})} + C, \end{aligned}$$

where C is a constant. Hence,

$$\int_0^{\pi/2} \frac{dx}{\cos \frac{x}{2} \cos^3 \frac{x}{6}} = 6 \int_0^{\pi/12} \frac{dt}{\cos 3t \cos^3 t} = \frac{8\sqrt{3}}{9} \left(2 \ln(\sqrt{3} + 1) - 10\sqrt{3} + \frac{33}{2} \right),$$

it follows that (5.5) holds. $\square$

The Schwab-Borchardt mean of two numbers $a \geq 0$ and $b > 0$, denoted by $SB(a, b)$, is defined as [2, Theorem 8.4], [6, 3, (2.3)]

$$SB(a, b) = \begin{cases} \frac{\sqrt{b^2 - a^2}}{\arccos(a/b)} & \text{if } a < b, \\ a & \text{if } a = b, \\ \frac{\sqrt{a^2 - b^2}}{\operatorname{arccosh}(a/b)} & \text{if } a > b. \end{cases}$$

The properties and certain inequalities involving Schwab-Borchardt mean can be found in [11], [12]. We now establish a new inequality for this mean.

Corollary 8. *For $t > 0$, we have*

$$(5.6) \quad (4t^2 - 1) SB(t, 1) \leq 3t^3.$$

Proof. For $t \in (0, 1)$, letting $\cos x = t$ in (4.2) we get

$$\frac{\sqrt{1 - t^2}}{\arccos t} (4t^2 - 1) < 3t^3.$$

For $t \in (1, \infty)$, we use Lin's inequality [9] of positive numbers $a, b > 0$ with $a \neq b$

$$\frac{a - b}{\ln a - \ln b} < \left(\frac{a^{1/3} + b^{1/3}}{2} \right)^3$$

to deduce

$$\frac{\sinh u}{u} < \cosh^3 \frac{u}{3}$$

by setting $u = \ln \sqrt{a/b}$. Letting $\cosh(u/3) = t$ and next using duplication formula for $\sinh$ function lead us to

$$\frac{\sqrt{t^2 - 1}}{\operatorname{arccosh} t} (4t^2 - 1) < 3t^3,$$

which proves the desired inequality. $\square$

The Seiffert's mean [16] of positive numbers $a, b > 0$ with $a \neq b$ is defined by

$$P = P(a, b) = \frac{a - b}{2 \arcsin \frac{a - b}{a + b}},$$

and let $A = A(a, b)$ and $G = G(a, b)$ denote the arithmetic mean and geometric mean. With $x = \arcsin \frac{a - b}{a + b} \in (0, \pi/2)$, we have

$$\frac{P}{A} = \frac{\sin x}{x}, \quad \frac{G}{A} = \cos x.$$

Thus Corollary 3 can be restated as follows.

Corollary 9. *For $a, b > 0$ with $a \neq b$, we have*

$$(5.7) \quad P < \frac{3}{4} \frac{(A + G)^2}{2G + A}.$$

REFERENCES

- [1] G. D. Anderson, M. K. Vamanamurthy AND M. Vuorinen, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, New York 1997.
- [2] M. Biernacki AND J. Krzyz, On the monotonicity of certain functionals in the theory of analytic functions, *Annales Universitatis Mariae Curie-Skłodowska* **9** (1995) 135–147.
- [3] D. M. Bradley, Representations of Catalan’s constant, 2001, available online at <http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.26.1879>.
- [4] J. M. Borwein AND P. B. Bowein, *Pi and the AGM: A Study in Analytic Number Theory and Computational Complexity*, John Wiley and Sons, New York, 1987.
- [5] E. Catalan, Recherches sur la constante G , et sur les intégrales eulériennes, *Mémoires de l’Académie impériale des sciences de Saint-Petersbourg, Ser.* **31** (7) (1883).
- [6] B. C. Carlson, Algorithms involving arithmetic and geometric means, *Amer. Math. Monthly* **78** (1971), 496–505.
- [7] Group of compilation, Handbook of Mathematics, Peoples’ Education Press, Beijing, China, 1979. (Chinese)
- [8] R. Klén, M. Visuri AND M. Vuorinen, On Jordan type inequalities for hyperbolic functions, *J. Inequal. Appl.* **2010** (2010), Art. ID 362548, 14 pages, doi:10.1155/2010/362548.
- [9] T. P. Lin, The power mean and the logarithmic mean, *Amer. Math. Monthly* **81** (1974), 879–883.
- [10] D. S. Mitrinović AND P. M. Vasić, *Analytic Inequalities*, Springer, New York, 1970.
- [11] E. Neuman AND J. Sándor, On the Schwab-Borchardt mean, *Math. Pannon.* **14** (2003), 253–266.
- [12] E. Neuman, Inequalities for the Schwab-Borchardt mean and their applications., *J. Math. Inequal.* in print.
- [13] F. Qi, Extensions and sharpenings of Jordan’s and Kober’s inequality, *Journal of Mathematics for Technology* **12** (4) (1996), 98–102. (Chinese)
- [14] F. Qi, D.-W. Niu AND B.-N. Guo, Refinements, generalizations, and applications of Jordan’s inequality and related problems, *J. Inequal. Appl.* **2009** (2009), Art. ID 271923, 52 pages.
- [15] F. Qi, L.-H. Cui AND S.-L. Xu, Some inequalities constructed by Tchebysheff’s integral inequality, *Math. Inequal. Appl.* **2** (4) (1999) 517–528.
- [16] H.-J. Seiffert, Werte zwischen dem geometrischen und dem arithmetischen Mittel zweier Zahlen, *Elem. Math.* **42** (1987), 105–107.
- [17] M. K. Vamanamurthy AND M. Vuorinen, Inequalities for means, *J. Math. Anal. Appl.* **183** (1994) 155–166.
- [18] Sh.-H. Wu AND L. Debnath, A new generalized and sharp version of Jordan’s inequality and its applications to the improvement of the Yang Le inequality, *Appl. Math. Letters* **19** (12) (2006), 1378–1384.
- [19] Sh.-H. Wu, Sharpness and generalization of Jordan’s inequality and its application, *Taiwanese J. Math.* **12** (2) (2008), 325–336.

SYSTEM DIVISION, ZHEJIANG PROVINCE ELECTRIC POWER TEST AND RESEARCH INSTITUTE,
 HANGZHOU, ZHEJIANG, CHINA, 310014
E-mail address: yzhkm@163.com